

Iterated Class Forcing

Sy D. Friedman*
M.I.T.

In this paper we develop the notion of “stratified” class forcing and show that this property both implies cofinality-preservation and is preserved by iterations with the appropriate support. Many Easton-style and Jensen-style forcings are stratified, as are some more exotic forcings obtained by mixing these types together (see Easton [70], section 36 of Jech [78], Beller-Jensen-Welch [82], Friedman [90]).

As a sample application, cofinalities are preserved by an iteration of length ORD where at even stages i , an Easton-style forcing adds a Cohen set to regular cardinals $\geq \text{card}(i)$, at odd stages $i + 1$ the class added at stage i is coded by a subset of the least infinite regular cardinal $\geq \text{card}(i)$ via the techniques of Friedman [93] or Friedman [94], and for any regular κ , $\{i | p(i) \text{ is nontrivial below } \kappa\}$ is a subset of κ^+ of size $< \kappa$, for each condition p in the iteration.

Jensen coding as in Beller-Jensen-Welch [82] is not stratified but obeys a related property, called Δ -stratification, which is also preserved by iterations with the appropriate (larger) support. As a sample application, the original form of Jensen coding can be used in the iteration of the preceding paragraph, provided the Cohen sets are added with full support at successor cardinals only and the condition stated at the end of that paragraph is imposed only at successor cardinals.

We now define stratification, in the language of Gödel-Bernays class theory.

Definition P (partially ordered by $\leq$) is *stratified* if there is a class A such that $V = L[A]$ has a $\langle V, A \rangle$ -definable well ordering and:

(a) P and $\leq$ are A -definable. A condition in P is a function p on an initial segment of $\text{Card} = \{0\} \cup \text{Infinite Cardinals}$, where if q extends p as a function, $q(\gamma) = \emptyset$ for all $\gamma \in \text{Dom}(q) - \text{Dom}(p)$, then we identify p with q . Also we require that $p(\kappa) = \emptyset$ for singular κ and the conditions with constant value $\emptyset$ are the weakest in P . Lastly, $\{p | p \restriction \kappa^+ \in H_{\kappa^+}\}$ is dense for each $\kappa \in \text{Card}$.

(b) (κ -Density Reduction) Let κ be regular and define $p \leq_\kappa q$ if $(p \leq q \text{ and } p \restriction \kappa^+ = q \restriction \kappa^+)$.

*Research supported by NSF Contract# 9205530-DMA.

$\kappa^+ = q \restriction \kappa^+$). Then $p \leq q \longrightarrow \exists r \leq_\kappa q \exists s \leq r (s \restriction [\kappa^+, \infty) = r \restriction [\kappa^+, \infty) \text{ and } s \leq p)$ and $p \leq q \longleftarrow \exists r \exists s (p \leq r, p \leq s \text{ and } r \restriction \kappa^+ = q \restriction \kappa^+, s \restriction [\kappa^+, \infty) = q \restriction [\kappa^+, \infty))$. If D is an A -definable dense class, p a condition then $\exists q \leq_\kappa p \exists d \subseteq D (\text{card}(d) \leq \kappa \text{ and every } r \leq q \text{ can be extended to } s \text{ such that } s \restriction [\kappa^+, \infty) = r \restriction [\kappa^+, \infty) \text{ and } s \text{ extends an element of } d)$.

(c) (κ -Definable Closure) For infinite regular κ there are Π_n^A operators $F_n(x, \kappa, p)$ for $0 < n \in \omega$ such that $F_n(x, \kappa, p) \leq_\kappa p$ for all p and whenever $p_0 \geq_\kappa p_1 \geq_\kappa \dots$ is a Π_n^A (in parameters from $\kappa \cup \{x\}$) sequence of length $\lambda \leq \kappa$ such that for each $i < \lambda$, $p_{i+1} \leq_{\kappa_i} F_n(x, \kappa_i, p'_i)$ for some $p'_i \leq_{\kappa_i} p_i$ and regular $\kappa_i \geq \kappa$ then there is $p \leq_\kappa p_i$ for all $i < \lambda$.

Theorem 1. *Suppose that P is stratified. Then P preserves ZFC (relative to the class A witnessing stratification), cofinalities and the GCH.*

Proof. Using κ -Definable Closure and κ -Density Reduction we get: If $\langle D_i | i < \kappa \rangle$ is a uniformly A -definable sequence of dense classes and $p \in P$ then there is $q \leq p$ and d of cardinality $\leq \kappa$ such that each $r \leq q$ is compatible with an element of $D_i \cap d$, for each i . This implies that the forcing relation for Δ_0^A sentences is A -definable and that A -replacement, cofinalities are preserved.

To show that GCH is preserved let $p \Vdash \tau \subseteq \kappa$, where κ is an infinite cardinal and define $D_i = \{q | q \text{ is incompatible with } p \text{ or } q \Vdash i \in \tau \text{ or } q \Vdash i \notin \tau\}$. Then D_i is dense and A -definable, uniformly in $i < \kappa$. First suppose that κ is regular. Then by κ -Density Reduction and κ -Definable Closure there is $q \leq p$ and d of size $\leq \kappa$ such that $r \in d \longrightarrow r \restriction [\kappa^+, \infty) = q \restriction [\kappa^+, \infty)$, $r \restriction \kappa^+ \in H_{\kappa^+}$ and $\{r \in d | r \Vdash i \in \tau \text{ or } r \Vdash i \notin \tau\}$ is predense below q for each $i < \kappa$. (X is predense below q if every extension of q is compatible with an element of X .) Thus in the generic extension each subset of κ is determined by a κ -sequence of size $\leq \kappa$ subsets of H_{κ^+} of the ground model, so the property $2^\kappa = \kappa^+$ is preserved.

When κ is singular the above argument can be repeated, using κ_i -Density Reduction and $\text{cof}(\kappa)$ -Definable closure, where $\langle \kappa_i | i < \text{cof}(\kappa) \rangle$ is a sequence of regular cardinals converging to κ . ⊥

To preserve stratification under iteration we must discuss strong witnesses and diagonal supports.

Definition P is stratified with *strong witness* A if in the definition of stratified,

$V = L[A]$ has a Δ_1^A -well ordering, $P, \leq, \text{Card}$ are Δ_1^A and there is a Σ_1^A function $f(x, \kappa, p) = (q, d)$ such that if x is an index for a Σ_1^A dense class D, p a condition and κ infinite and regular then $q \leq_\kappa p$, $\text{card}(d) \leq \kappa, d \subseteq D$ and every $r \leq q$ can be extended to s such that $s \restriction [\kappa^+, \infty) = r \restriction [\kappa^+, \infty)$ and s extends an element of d .

Proposition 2. *If P is stratified then P has a strong witness.*

Proof. Suppose A witnesses that P is stratified. Then we can choose $A^* = \Sigma_N^A$ satisfaction, for a large N so that V has a $\Delta_1^{A^*}$ well ordering and $P, \leq, \text{Card}, \{(q, d, \kappa) \mid \text{card}(d) \leq \kappa \text{ and every } r \leq q \text{ can be extended to } s \text{ such that } r \restriction [\kappa^+, \infty) = s \restriction [\kappa^+, \infty) \text{ and } s \text{ extends an element of } d\}$ are $\Delta_1^{A^*}$. Then the desired Σ_1^A function $f(x, \kappa, p)$ exists. Finally define the $\Pi_n^{A^*}$ operator $F_n^*(x, \kappa, p)$ to be $F_{N+n}(x, \kappa, p)$ where $\langle F_m(x, \kappa, p) \mid 0 < m \in \omega \rangle$ comes from A . $\dashv$

Strong witnesses help us control the definability of the forcing relation.

Theorem 3. *Suppose that A is a strong witness to the stratification of P . Then the forcing relation for P restricted to Σ_1^A sentences is densely- Σ_1^A : there is a Σ_1^A relation $p \Vdash^* \varphi$ ($p \in P, \varphi \in \Sigma_1^A$) such that $p \Vdash^* \varphi \longrightarrow p \Vdash \varphi$ and $p \Vdash \varphi \longrightarrow \exists q \leq p, q \Vdash^* \varphi$.*

Proof. It suffices to prove this for $\Delta_0 \varphi$, by looking at witnesses. We show by a Σ_1^A induction on φ that given p we can (in a Σ_1^A way) find $q \leq p$ and $i \in \{0, 1\}$ such that either $q \Vdash \varphi, i = 0$ or $q \Vdash \neg \varphi, i = 1$. This will prove the Theorem since we can then take $q \Vdash^* \varphi \longleftrightarrow$ For some $p, (q, 0)$ arises from p, φ as above.

The interesting case of the induction is the bounded quantifier: Suppose φ is $\forall x \in \sigma \psi(x)$ where σ is a term of rank α . By induction we can effectively extend p to decide any instance $\psi(\tau)$, $\text{rank}(\tau) < \alpha$. If one of these extensions q_τ forces $\tau \in \sigma \wedge \neg \psi(\tau)$ then we can take the desired q to be $q_\tau \Vdash \neg \varphi$. Otherwise, we can build uniformly Σ_1^A dense classes D_τ , $\text{rank}(\tau) < \alpha$ of conditions forcing $\tau \in \sigma \longrightarrow \psi(\tau)$. As A is a *strong* witness we can effectively find d and $q \leq p$ such that each $D_\tau \cap d$ is predense below q , where d is a set. But this q forces $\forall x \in \sigma \psi(x) = \varphi$. $\dashv$

We are ready to discuss stratified iterations.

Definition $\langle P_i | i < \alpha \rangle$ (where $P_{i+1} = P_i * \mathbb{Q}_i$, $P_\lambda \subseteq \text{Inverse Limit } \langle P_i | i < \lambda \rangle$ for limit λ) is a *stratified iteration* if for some class $A \subseteq \text{ORD}$, A strongly witnesses that P_0 is stratified, for each $i + 1 < \alpha$, $\emptyset_i \Vdash_i \mathbb{Q}_i$ is stratified with strong witness $\langle A, G_i \rangle$ via some F_n^i, f^i ($\emptyset_i =$ weakest condition in $P_i, \Vdash_i =$ forcing for $P_i, G_i =$ generic for P_i) and $\mathbb{Q}_i, f^i$ are $\tilde{\Delta}_1^{A, G_i}$ uniformly in $i < \alpha$, F_n^i is $\tilde{\Pi}_n^{A, G_i}$ uniformly in $i < \alpha$, for each $n > 0$. Such an iteration has *short diagonal supports* if for $p \in P_j$ and infinite regular κ , $\{i | i < j \text{ and } p \restriction i \not\Vdash_i \forall \gamma < \kappa^+, p(i)(\gamma) = \emptyset\}$ is a subset of κ^+ of size $< \kappa$ (and this is the only restriction on supports).

Stratification Theorem. *Suppose $\langle P_i | i < \alpha \rangle$ is a stratified iteration with short diagonal supports and GCH holds. Then P_α is isomorphic to a stratified forcing (definably relative to a class A witnessing stratification).*

Proof. First we note that in the definition of stratified, we may assume one further condition about the operators $F_n(x, \kappa, p) : \kappa_1 \leq \kappa_2$ both regular, $p(\gamma) = \emptyset$ for all $\gamma < \kappa_2 \longrightarrow F_n(x, \kappa_1, p) = F_n(x, \kappa_2, p)$. For, we may achieve this property by redefining F_n to be $F_n^*(x, \kappa, p) = F_n(x, \kappa(p), p)$ where $\kappa(p) =$ least γ such that $p(\gamma) \neq \emptyset$, if $\kappa \leq \kappa(p)$ and $F_n^*(x, \kappa, p) = F_n(x, \kappa, p)$ otherwise.

We prove the Main Theorem by induction on α , maintaining the coherence property that the isomorphism of P_α with a stratified forcing P_α^* extend the (inductively produced) isomorphism of P_β with the stratified forcing P_β^* for $\beta < \alpha$, viewing P_β as a subforcing of P_α in the natural way (and P_β^* as a subforcing of P_α^* in a natural way that will be evident from the construction). Also if A is our given witness to the stratification of the iteration then A will serve as a strong witness to the stratification of each P_α^* .

The result is vacuous for $\alpha = 0$ or 1 . Suppose that $\alpha = \beta + 1$ is a successor ordinal > 1 . By induction P_β is isomorphic to a stratified forcing P_β^* and let $\leq_\kappa^\beta, F_n^\beta(x, \kappa, p^\beta)$ come from the stratification of P_β^* . Also $\emptyset_\beta \Vdash_\beta \mathbb{Q}_\beta^*$ is stratified ($\emptyset_\beta =$ weakest condition of $P_\beta^*, \Vdash_\beta =$ forcing for $P_\beta^*, \mathbb{Q}_\beta^* =$ the P_β^* -name for $\mathbb{Q}^\beta$) and let $\leq_\kappa, F_n(x, \kappa, q)$ result from this.

By Theorem 3, $\Vdash_\beta$ is densely $\tilde{\Sigma}_1^A$ when restricted to $\tilde{\Sigma}_1^A$ sentences. By replacing A by $\tilde{\Sigma}_1^A$ -Satisfaction, we may assume in fact that $\Vdash_\beta$ is $\tilde{\Sigma}_1^A$ for $\tilde{\Sigma}_1$ sentences. It follows that $\Vdash_\beta$ is $\tilde{\Sigma}_n^A$ when restricted to $\tilde{\Sigma}_n$ sentences.

Now we define P_α^* to essentially consist of all functions f on an initial segment of

Card such that for some p^β in P_β^* and some $q, p^\beta \Vdash_\beta q \in \mathbb{Q}_\beta^*$ and for all $\kappa \in \text{Dom}(f)$, $f(\kappa) = \langle p^\beta(\kappa), q(\kappa) \rangle$ where $q(\kappa)$ is the canonical term denoting the result of applying the function denoted by q to κ . However we must make two small modifications: insist that if $p^\beta(\kappa) = \emptyset$ and $p^\beta \Vdash_\beta q(\kappa) = \emptyset$ or undefined then $f(\kappa) = \emptyset$ (instead of $\langle \emptyset, \text{a term for } \emptyset \rangle$); also insist that $\text{Dom}(f)$ contains $\text{Dom}(p^\beta)$ and $\text{rank}(q) < \cup \text{Dom}(f)$, so that $\emptyset_\beta \Vdash_\beta \text{Dom}(q) \subseteq \text{Dom}(f)$. Then clearly P_α^* is isomorphic to P_α when P_α^* is ordered in the natural way (by ordering the corresponding pairs $\langle p^\beta, q \rangle$ in $P_\beta^* * \mathbb{Q}_\beta^*$). It is easy to verify condition (a) and the first part of (b) in the definition of stratification.

Next we demonstrate κ -Density Reduction for P_α^* . For notational purposes we think of a condition in P_α^* as an element of $P_\beta^* * \mathbb{Q}_\beta^*$ (isomorphic to P_α^*). Suppose $D \subseteq P_\alpha^*$ is dense and A -definable and $(p^\beta, q) \in P_\alpha^*$. Consider $D^{G_\beta^*} = \{q_0 \in \mathbb{Q}_\beta^* \mid (p_0^\beta, q_0) \in D \text{ for some } p_0^\beta \in G_\beta^*\}$ where G_β^* denotes the P_β^* -generic. Then $D^{G_\beta^*} \subseteq \mathbb{Q}_\beta^*$ is forced by $\emptyset_\beta \in P_\beta^*$ to be dense. So by κ -Density Reduction for $\mathbb{Q}_\beta^*$, $\emptyset_\beta^*$ also forces that $\{q_0 \in \mathbb{Q}_\beta^* \mid q_0 \text{ reduces } D^{G_\beta^*} \text{ below } \kappa^+, \text{ to size } \leq \kappa\}$ is $\leq_\kappa$ -dense on $\mathbb{Q}_\beta^*$ (i.e., that for any q_0 there is $q_1 \leq_\kappa q_0$ such that for some $d \subseteq D^{G_\beta^*}$ of size $\leq \kappa$, every $r \leq q_1$ can be extended to s such that s, r agree $\geq \kappa^+$ and s extends an element of d). Thus $D_\beta = \{p_0^\beta \mid \text{For some } q_0, d_0, p_0^\beta \Vdash_\beta q \notin \mathbb{Q}_\beta^* \text{ or } q_0 \leq_\kappa q \text{ reduces } D^{G_\beta^*} \text{ below } \kappa, \text{ to } d_0, \text{ card}(d_0) \leq \kappa\}$ is dense on P_β^* . Let $p_0^\beta \leq_\kappa^\beta p^\beta$ reduce D_β below κ^+ , to size $\leq \kappa$, by κ -Density Reduction for P_β^* .

Then we can form terms q_0, d_0 such that $p_0^\beta \Vdash_\beta q_0 \leq_\kappa q$, q_0 reduces $D^{G_\beta^*}$ below κ^+ , to d_0 of size $\leq \kappa$. For each $i < \kappa$ it is dense below p_0^β to force some $p_0^\beta(i) \in G_\beta^*$, $(p_0^\beta(i), d_0(i)) \in D$, where $d_0(i) = i^{\text{th}}$ element of d_0 . Finally, by κ -Density Reduction and κ -Definable Closure for P_β^* , we can assume that p_0^β reduces all of these dense sets below κ^+ , to size $\leq \kappa$ and hence (p_0^β, q_0) reduces D below κ^+ , to size $\leq \kappa$.

To complete the successor case we need to define the operators $F_n^{\beta+1}(x, \kappa, (p^\beta, q))$ and verify condition (c) in the definition of stratified. We set $F_n^{\beta+1}(x, \kappa, (p^\beta, q)) = \text{“least” } (p_0^\beta, q_0) \text{ such that } p_0^\beta \leq_\kappa^\beta F_n^\beta(x, \kappa, p^\beta) \text{ and } p_0^\beta \Vdash_\beta F_n(x, \kappa, q) = q_0$. Note that the property of (p_0, q_0) stated here is Σ_n^A (in the other parameters), so we take “least” in the sense of Σ_n^A -uniformization, so that $F_n^{\beta+1}(x, \kappa, p^{\beta+1})$ is Σ_n^A . Of course we must show that such a (p_0^β, q_0) exists. Note that it is a dense property of p_0^β to force a value for $F_n(x, \kappa, q)$. By κ -Density Reduction there is p_0^β reducing this

dense property to a set, with $p_0^\beta \leq_\kappa^\beta F_n^\beta(x, \kappa, p^\beta)$. Thus we can form a term q_0 such that $p_0^\beta \Vdash_\beta F_n(x, \kappa, q) = q_0$.

The κ -Definable Closure of $P_{\beta+1}^*$ follows from the κ -Definable Closure of P_β^* (relative to A) and the κ -Definable Closure of $\mathbb{Q}_\beta^*$ (relative to $\langle A, G_\beta^* \rangle$). Also $\leq_\kappa^{\beta+1}$ is Δ_1^A -definable, uniformly in κ , using the facts that $\leq_\kappa^\beta$ is uniformly Δ_1^A , $\leq_\kappa$ is uniformly Δ_1^{A, G_β} and the fact that $\Vdash_\beta$ is Σ_1^A when restricted to Σ_1 sentences.

Now we turn to the case where α is a limit ordinal. We take P_α^* to consist of all functions f on an initial segment of Card such that for some $\langle f_\beta | \beta < \alpha \rangle$ in the inverse limit of $\langle P_\beta^* | \beta < \alpha \rangle$ with short diagonal supports, $f(\kappa) = \langle f_\beta(\kappa) | \beta < \alpha \rangle$ for all κ in $\text{Dom}(f)$; we also require that $\text{Dom}(f) \supseteq \text{Dom}(f_\beta)$ for each $\beta < \alpha$ and modify $f(\kappa)$ to be $\emptyset$ if $f_\beta(\kappa) = \emptyset$ for all $\beta < \alpha$. The f 's are ordered by ordering the corresponding $\langle f_\beta | \beta < \alpha \rangle$'s.

We must show that $\{f \in P_\alpha^* | f \restriction \kappa^+ \in H_{\kappa^+}\}$ is dense for each κ . We actually show a bit more, for the purpose of carrying out an inductive argument: if $\gamma < \kappa$ belong to Card , γ regular then $\{f \in P_\alpha^* | f \restriction [\gamma, \kappa] \in H_{\kappa^+}\}$ is $\leq_\gamma$ -dense (any f can be $\leq_\gamma$ -extended into this set; for $\gamma = 0$ take $\leq_\gamma = \leq$.) Note that this stronger version follows from the weaker one, given γ -Density Reduction, so we may inductively assume that it holds for $P_\beta^*, \beta < \alpha$. Now we induct on κ : using short diagonal supports, we may assume that $\text{cof}(\alpha) < \kappa$ as otherwise our given f has the property that for some $\beta_0 < \alpha$, f_β is the $\emptyset$ -function below κ^+ for all $\beta_0 \leq \beta < \alpha$ (where f comes from $\langle f_\beta | \beta < \alpha \rangle$) and so we can apply induction at β_0 . By induction on κ we can first extend f to guarantee that $f \restriction [\gamma, \text{cof}(\alpha))$ belongs to $H_{\text{cof}(\alpha)^+}$. So we may assume that $\gamma \geq \text{cof}(\alpha)$. Now, choose a cofinal $\text{cof}(\alpha)$ -sequence $\alpha_0 < \alpha_1 < \dots$ below α and successively extend $f = f_0$ to $f_1, f_2, \dots$ in $\text{cof}(\alpha)$ steps so that $(f_{i+1} \restriction \alpha_i)$ on $[\gamma, \kappa]$ belongs to H_{κ^+} and $f_{i+1} \restriction \alpha_j \geq_\gamma F_1^{\alpha_j}(x, \gamma, f_i \restriction \alpha_j)$ for all $j \leq i$, where $F_1^{\alpha_i}$ comes from Definable Closure for P_{α_i} and $x = \langle f, \gamma, \kappa, \langle \alpha_i | i < \text{cof} \alpha \rangle \rangle$. (We abuse notation slightly; $f_i \restriction \alpha_i$ actually should be the function $g(\delta) = f_i(\delta) \restriction \alpha_i$.) Note that a simple construction using the $F_1^{\alpha_j}$'s shows that f_{i+1} as above does exist. So we get that f_λ is a condition for limit λ and $f_{\text{cof}(\alpha)}$ is as desired.

If $\text{cof}(\alpha) \leq \kappa$ we define $F_n^\alpha(x, \kappa, p)$ to be the least $q \leq_\kappa p$ such that $q \restriction \alpha_i \leq_\kappa F_n^{\alpha_i}(x, \kappa, p \restriction \alpha_i)$ for each α_i in a fixed $\text{cof}(\alpha)$ -sequence cofinal in α . If $\kappa < \text{cof}(\alpha) < \alpha$ and $\text{cof}(\alpha)$ is not the successor of a regular cardinal then we obtain $F_n^\alpha(x, \kappa, p)$ by first choosing $q_0 \leq_{\text{cof}(\alpha)} p$ so that $q_0 \restriction \alpha_i \leq_{\text{cof}(\alpha)} F_n^{\alpha_i}(x, \text{cof}(\alpha), p \restriction \alpha_i)$ for

each α_i and then $q_1 \leq_\kappa q_0$ so that $q_1 \restriction \text{cof}(\alpha) \leq_\kappa F_n^{\text{cof}(\alpha)}(x, \kappa, q_0 \restriction \text{cof}(\alpha))$. If $\kappa \leq \lambda < \lambda^+ = \text{cof}(\alpha) < \alpha$ with λ regular then we choose q_0, q_1 as above and then $q_2 \leq_\lambda q_1$ so that $q_2 \restriction \beta \leq_\lambda F_n^\beta(x, \lambda, q_1 \restriction \beta)$ for each β such that $\text{cof}(\alpha) \leq \beta$ and $q_1 \restriction \beta \not\leq_\beta q_1(\beta)$ is the $\emptyset$ -function below $\text{cof}(\alpha)$. Finally if $\kappa < \alpha$ and α is regular then choose q_0 as above ($\alpha_i = i$) and then $q_1 \leq_\kappa q_0$ so that $q_1 \restriction \beta \leq_\kappa F_n^\beta(x, \kappa, q_0 \restriction \beta)$ where $\beta < \alpha$ is least so that $\beta \leq \beta' \longrightarrow q_0 \restriction \beta' \Vdash_{\beta'} q_0(\beta')$ is the $\emptyset$ -function below α^+ . Our construction guarantees that if $q = F_n^\alpha(x, \kappa, p)$ and $\beta < \alpha$ then for some $\kappa' \geq \kappa$ and $q', q \restriction \beta \leq_\kappa q' \leq_{\kappa'} F_n^\beta(x, \kappa', p')$ for some $p' \leq_\kappa p \restriction \beta$. The latter is used to verify κ -Definable Closure when α is regular. (The other cases are straightforward, using our extra hypothesis about the F_n 's stated at the start of the proof.)

Finally we must establish the second part of κ -Density Reduction for P_α^* . (The first part is easy if $\text{cof}(\alpha) \leq \kappa$ and otherwise follows inductively.) First suppose that $\alpha < \kappa^+$ and choose an increasing cofinal $\alpha_0 < \alpha_1 < \dots$ of ordertype $\text{cof}(\alpha)$. Given $p \in P_\alpha^*$ and A -definable open dense D , use the $F_n^{\alpha_i}$ functions to successively extend $p \restriction \alpha_i$ producing $q \leq_\kappa p$ such that for each $i < \text{cof } \alpha$, $q \restriction \alpha_i$ reduces D^{α_i} below κ^+ to size $\leq \kappa$, where $D^{\alpha_i} = \{r \restriction \alpha_i \mid r \in D\}$. Now successively $\leq_\kappa$ -extend $q = q_0$ to $q_1, q_2, \dots$ so that for each γ there is $x_{\gamma+1}$ defined on $\text{Card} \cap \kappa^+$ so that $x_{\gamma+1} \cup q_{\gamma+1}$ is an element of D yet is incompatible with each $x_{\gamma'+1} \cup q_{\gamma'+1}$ for $\gamma' < \gamma$. But for each i there must be a stage $\gamma_i < \kappa^+$ such that for $\gamma \geq \gamma_i$, $(x_{\gamma+1} \cup q_{\gamma+1}) \restriction \alpha_i$ is compatible with some $(x_{\gamma'+1} \cup q_{\gamma'+1}) \restriction \alpha_i$ where $\gamma' < \gamma_i$, since $q \restriction \alpha_i$ reduces D^{α_i} below κ^+ to size $\leq \kappa$. Let $\gamma = \cup \{\gamma_i \mid i < \text{cof } \alpha\} < \kappa^+$. Then $q_{\gamma+1}$ is undefined so some $q_{\gamma'}, \gamma' < \kappa^+$ reduces D below κ^+ , to size $\leq \kappa$.

Now suppose that $\alpha \geq \kappa^+$. We may assume that $\alpha = \kappa^+$ as short diagonal supports requires that $p \in P_\alpha^*$ are trivial below κ^+ on all but fewer than κ coordinates, all below κ^+ . But note that we can assume that conditions in D when restricted to $\text{Card} \cap \kappa^+$ belong to H_{κ^+} and therefore can choose $q \leq_\kappa p$ and $\alpha_o < \kappa^+$ of cofinality κ such that the conditions in D which are trivial below κ^+ on coordinates $\geq \alpha_o$ form a set predense below q . If we extend q to $q_0 \leq_\kappa q$ reducing D^{α_o} below κ^+ to size $\leq \kappa$, then q_0 in fact reduces D below κ^+ to size $\leq \kappa$. $\dashv$

There are some important examples of cofinality-preserving class forcings that are not stratified. Instead they may obey Δ -stratification, which we now consider.

Definition P is Δ -stratified if it obeys the definition of stratified where (b), (c)

are restricted to successor cardinals and in addition: whenever $0 < n \in \omega$ and κ is inaccessible, $p_0 \geq p_1 \geq \dots$ a Π_n^A (in parameters from $\lambda \cup \{x\}$) sequence of length $\lambda \leq \kappa$ and for each $i < \lambda$, $p_{i+1} \leq_{\kappa_i} F_n(x, \kappa_i, p'_i)$ for some $p'_i \leq_{\kappa_i} p_i$ and regular $\kappa_i \geq \aleph_{i+1}$, there is $p \leq_{\aleph_{i+1}} p_i$ for each i .

A is a *strong witness* to the Δ -stratification of P if it obeys the definition of strong witness to stratification when restricted to successor cardinals. A Δ -*stratified iteration* is just like a stratified iteration but with stratified replaced by Δ -stratified everywhere. Such an iteration $\langle P_i | i < \alpha \rangle$ has *long diagonal supports* if for $p \in P_j$ and successor cardinals κ , $\{i < j | p \restriction i \not\vdash \forall \gamma \leq \kappa, p(i)(\gamma) = \emptyset\}$ is a subset of κ^+ of size $< \kappa$ and for inaccessible $\kappa \leq j$, $\{\bar{\kappa} < \kappa | \text{For some } \bar{\kappa} \leq j' < j, p \restriction j' \not\vdash_{j'} p(j') \text{ is } \emptyset \text{ at } \bar{\kappa}\}$ is nonstationary in κ (and these are the only support restrictions).

Theorem 4. *Suppose that P is Δ -stratified. Then P preserves ZFC (relative to the class A witnessing Δ -stratification), cofinalities and the GCH.*

Proof. As in the proof of Theorem 1, using Δ -stratification at κ and stratification at $\bar{\kappa}^+ < \kappa$, when κ is inaccessible. $\dashv$

Δ -Stratification Theorem. *Suppose $\langle P_i | i < \alpha \rangle$ is a Δ -stratified iteration with long diagonal supports and GCH holds. Then P_α is isomorphic to a Δ -stratified forcing (definably relative to a class A witnessing Δ -stratification).*

Proof. We follow the proof of the Stratification Theorem. Note that Theorem 3 still applies since its proof only uses that conditions (b), (c) hold at cofinally many regular cardinals. We proceed by induction on α . For successor α our earlier proof still shows that (b), (c) hold at successor cardinals. For Δ -stratification at an inaccessible κ , use Δ -stratification for $P_\beta^*(\alpha = \beta + 1), \emptyset_\beta \Vdash_\beta \Delta$ -stratification for $\mathbb{Q}_\beta^*$ to obtain $\bar{p} \restriction \beta \leq_{\aleph_{i+1}} p_i \restriction \beta$ for each i , $\bar{p} \restriction \beta \Vdash_\beta$ there is $\bar{p}(\beta) \leq_{\aleph_{i+1}} p_i(\beta)$ for each i and then $\leq_{\kappa^+}$ extend $\bar{p} \restriction \beta$ to $p \restriction \beta$ so that for some term $p(\beta)$, $p \restriction \beta \Vdash_\beta p(\beta) \leq_{\aleph_{i+1}} p_i(\beta)$ for each i , using κ^+ -Density Reduction. Then $p \leq_{\aleph_{i+1}} p_i$ for each i is as desired.

When α is a limit ordinal we define P_α^* as before and first show that $\{f \in P_\alpha^* | f \restriction [\gamma, \kappa] \in H_{\kappa^+}\}$ is $\leq_\gamma$ -dense for each successor $\gamma < \kappa$, γ and κ in Card. We do this by induction on κ , noting that we may assume it holds for $P_\beta^*, \beta < \alpha$. Using (long) diagonal supports we may assume that either $\alpha = \kappa$ is inaccessible

or $\text{cof}(\alpha) < \kappa$. If $\text{cof}(\alpha)$ is a successor or $\text{cof}(\alpha)^+ < \kappa$ then the old argument can be applied, using $\text{cof}(\alpha)$ -Definable Closure or $\text{cof}(\alpha)^+$ -Definable Closure applied to $P_{\alpha_i}^*$, $\alpha_i < \alpha$. So we may assume that either $\alpha = \kappa$ is inaccessible or $\text{cof}(\alpha)^+ = \kappa$ where $\text{cof}(\alpha)$ is inaccessible. In the latter case we choose a cofinal $\text{cof}(\alpha)$ sequence $\alpha_0 < \alpha_1 \dots$ and successively $\leq_\gamma$ -extend our given $f = f_0$ to $f_1, f_2, \dots$ in $\text{cof}(\alpha)$ steps so that $(f_{i+1} \upharpoonright \alpha_i) (\kappa) \in H_{\kappa^+}$ and $f_{i+1} \upharpoonright \alpha_j \geq_{\aleph_{j+1}} F_1^{\alpha_j}(x, \aleph_{j+1} \cup \gamma, f_i \upharpoonright \alpha_j)$ for all $j \leq i$, $x = \langle f, \gamma, \kappa, \langle \alpha_i | i < \text{cof}(\alpha) \rangle \rangle$. Note that by induction we may extend $g = f_{\text{cof}(\alpha)}$ so that $g \upharpoonright [\gamma, \text{cof}(\alpha)] \in H_\kappa$, as desired. Finally if $\alpha = \kappa$ is inaccessible use Definable Closure to successively $\leq_\gamma$ -extend $f = f_0$ to $f_1, f_2, \dots$ in κ steps choosing a continuous cofinal $\kappa_0 < \kappa_1 < \dots$ below κ such that $f_{i+1} \upharpoonright (\kappa_i, \kappa_{i+1})$ belongs to $H_{\kappa_{i+1}^+}$ and $f_{i+1}(\kappa_i) = \emptyset$ for all i , using the fact that $\{\gamma < \kappa | f(\gamma) \neq \emptyset\}$ is nonstationary in κ . Then f_κ is as desired.

If $\text{cof}(\alpha) \leq \kappa$ or α is a successor cardinal nor $\text{cof}(\alpha)$ is neither inaccessible nor the successor of an inaccessible then we define $F_n^\alpha(x, \kappa, p)$ as in the stratified case. If $\text{cof}(\alpha) > \kappa$ is inaccessible then let $\alpha_0 < \alpha_1 < \dots$ be a cofinal $\text{cof}(\alpha)$ -sequence so that $\alpha_j \geq \aleph_{j+2} \cup \kappa$ for each $j < \text{cof}(\alpha)$ and let $F_n^\alpha(x, \kappa, p)$ be a lower bound of $p = p_0, p_1, \dots$ where p_{j+1} is least so that $p_{j+1} \upharpoonright \alpha_{j'} \leq_{\kappa \cup \aleph_{j+1}} F_n^{\alpha_{j'}}(x, \kappa \cup \aleph_{j+1}, p_j \upharpoonright \alpha_{j'})$ for all $j' \leq j$. If $\kappa \leq \lambda < \lambda^+ = \text{cof}(\alpha) < \alpha$, λ inaccessible then similarly modify the earlier definition of q_2 , enumerating the relevant β 's in λ steps.

κ -Density Reduction for successor κ follows just as in the stratified case. Δ -stratification also follows as our construction implies that if $p_{i+1} \leq_{\aleph_{i+1}} F_n^\alpha(x, \aleph_{i+1}, p_i)$ for $i < \kappa$ (κ inaccessible) then for cofinally many $\alpha' < \alpha$, $p_{i+1} \upharpoonright \alpha' \leq_{\aleph_{i'+1}} F_n^{\alpha'}(x, \aleph_{i'+1}, p_i \upharpoonright \alpha')$ for each i (and some $i' \geq i$ depending on α', i). Also if $\beta < \alpha$ and $p \leq_\kappa q$ in $P_{\beta+1}^*$, p at $\beta = q$ at β then $F_n^{\beta+1}(x, \kappa, p)$ at β equals $F_n^{\beta+1}(x, \kappa, q)$ at β . So given $p_0, p_1, \dots$ of length λ as in the hypothesis of Δ -stratification at κ for P_α^* , we can obtain the desired lower bound p by choosing $q \upharpoonright \beta + 1$ to be a lower bound for $\langle p \upharpoonright \beta \cup \{\langle \beta, p_i \text{ at } \beta \rangle | i < \lambda \rangle$ and taking $p(\beta) = q(\beta)$. $\dashv$

Examples

(a) Jensen coding (Beller-Jensen-Welch [82]) is equivalent to a Δ -stratified forcing. It is dense to have $p(0) \neq \emptyset$ and restricted to such conditions (together with the $\emptyset$ conditions) condition (a) is satisfied. (We must reindex though: $p(\kappa) = p(\kappa^+)$ in Jensen's sense.) The first part of (b) is clear at successor κ and the second part

is one of Jensen's key lemmas. For (c) we take $F_n(x, \kappa, p)$ to be the least $q \leq_\kappa p$ such that for $\lambda \leq \gamma \in \text{Dom}(p)$, $\gamma \in \Sigma_{n-1}^A \text{Hull}(\gamma \cup \{x\})$, $(q)_\gamma$ meets all predense D on P_γ in $\Sigma_{n-1}^A \text{Hull}(\gamma \cup \{x, p\})$. Jensen's lemmas show that such a q exists and that (c) is satisfied (one can assume that all the κ_i 's are equal by looking at their $\liminf$). The extra Δ -stratification condition also follows from Jensen's work.

(b) The modification of Jensen coding in Friedman [93] is equivalent to a forcing that is both stratified and Δ -stratified. It is densely embeddable in the forcing defined in the same way (after reindexing) but where at limit cardinals κ , we allow $p \restriction \kappa$ to code only an initial segment of p_κ (and belong to the coding structure for that initial segment). This allows one to prove (c) at inaccessibles. The thinning that was done there in the limit coding enables one to prove (b) at inaccessibles.

(c) The modification of Jensen coding in Friedman [94] is stratified. The proof of (b) at inaccessibles uses the fact that conditions have Easton domains.

(d) Easton forcing (see Easton [70]) where a Cohen set is added to each regular cardinal via an Easton product is stratified. (Take $F_n(x, \kappa, p) = p$.) If, instead, the full product is used but only at successor cardinals (no restriction on the domains of conditions) then Δ -stratification is obtained (but (b) will hold only at successors). Without the restriction to successor cardinals one has a "hybrid" forcing that is neither stratified nor Δ -stratified. Iterating it would require use of "mixed support."

(e) The forcing of Friedman [90] is a mixture of Jensen-style and Easton-style forcing. It is equivalent to a stratified forcing, provided one of the stratified modifications of Jensen coding (see (b), (c) above) is used.

(f) Backwards Easton forcings with Easton support (see Jech [78], section 36) are stratified provided at regular κ one uses a κ^+ -CC forcing of size $\leq \kappa^+$.

REFERENCES

- [82] Beller-Jensen-Welch, **Coding The Universe**, Cambridge University Press.
- [70] Easton, *Powers of Regular Cardinals*, Annals of Mathematical Logic.
- [90] Friedman, *The Π_2^1 -Singleton Conjecture*, Journal of the AMS.
- [93] Friedman, *A Simpler Proof of Jensen's Coding Theorem*, To appear Annals of Pure and Applied Logic.
- [94] Friedman, *A Short Proof of Jensen's Coding Theorem Assuming Not $0^\#$* , To appear.
- [78] Jech, **Set Theory**.